


# Superspace coinvariants and hyperplane arrangements

arXiv:2404.17919



by Trevor K. Karn (U. Minnesota)  
(joint with Robert Angarone, Patricia Commins, Satoshi Murai,  
and Brendon Rhoades)  
on Monday, 4 November, 2024

Main problem:

find a linear basis for the algebra  $SR_n$ .

Approach:

$ST$  algebras of southwest arrangements

What is  $SR_n$ ?

What are SW arrangements and  $ST$  algebras?

Proof ideas

What is  $SR_n$ ?

The symmetric group  $\mathfrak{S}_n$  acts on

$$\mathbb{C}[\mathbf{x}] = \mathbb{C}[x_1, x_2, \dots, x_n].$$

Let

$$p_k = x_1^k + x_2^k + \dots + x_n^k.$$

and let

$$I_n^+ = (p_1, p_2, \dots, p_n)$$

Definition

$R_n = \mathbb{C}[\mathbf{x}]/I_n^+$  is the coinvariant ring.

## » Example

Let  $n = 3$ ,  $\mathbb{C}[\underline{x}] = \mathbb{C}[x, y, z]$ ,

$$p_1 = x + y + z$$

$$p_2 = x^2 + y^2 + z^2$$

$$p_3 = x^3 + y^3 + z^3$$

so

$$x^4 + xy^3 + xz^3 \in I_3^+$$

and

$$x^2 + y^2 \notin I_3^+$$

In  $\mathbb{C}[\underline{x}]/I_3^+$ ,

$$x + y + z \equiv 0$$

and

$$x^2 + y^2 + z^2 \equiv (-y - z)^2 + y^2 + z^2.$$

so

$$y^2 \equiv -yz - z^2.$$

Similar computation shows that

$$z^3 \equiv 0.$$


Observation:  $x_i^j$  can always be rewritten “nicely”

Definition

The staircase monomials are  $x^\alpha$  where  $\alpha_i < i$ .



but not  $x_2^2 \longleftrightarrow$



Theorem (E. Artin)

The staircase monomials are a basis for  $R_n$ .



# What is $SR_n$ ?

Superspace is

$$\mathbb{C}[\underline{x}, \underline{\theta}] = \mathbb{C}[x_1, x_2, \dots, x_n, \theta_1, \theta_2, \dots, \theta_n]$$

where  $\theta_i\theta_j = -\theta_j\theta_i$ . Write<sup>†</sup>

$$\underline{\theta}^J = \prod_{j \in J} \theta_j.$$

$\mathfrak{S}_n$  acts simultaneously, so  $sp_k = x_1^k\theta_1 + x_2^k\theta_2 + \dots + x_n^k\theta_n$  is symmetric.

Let  $SI_n^+ = (p_1, \dots, p_n, sp_0, \dots, sp_{n-1})$ .

Definition

The superspace coinvariant ring is

$$SR_n = \mathbb{C}[\underline{x}, \underline{\theta}] / SI_n^+.$$

Sagan and Swanson [SS24] introduced

$$\mathcal{M} = \bigcup_{J \subseteq [n]} \{\underline{x}^\alpha \underline{\theta}^J : \alpha < (J\text{-staircase})\}$$

Definition

Let  $J \subseteq [n]$ . A  $J$ -staircase is  $(\text{st}(J)_1, \text{st}(J)_2, \dots, \text{st}(J)_n)$  where

$$\text{st}(J)_1 = \begin{cases} 0 & 1 \in J \\ 1 & 1 \notin J \end{cases}$$

and

$$\text{st}(J)_i = \begin{cases} \text{st}(J)_{i-1} & i \in J \\ \text{st}(J)_{i-1} + 1 & i \notin J \end{cases}$$

## » Example

Let  $J = \{2, 4, 5\} \subseteq [6]$ . Then the  $J$ -staircase is  $(1, 1, 2, 2, 2, 3)$ .



So  $\mathcal{M}$  contains monomials

$$x_3 x_6 \theta_2 \theta_4 \theta_5$$

and

$$x_4 x_6^2 \theta_2 \theta_4 \theta_5$$

but not

$$x_5^2 \theta_2 \theta_4 \theta_5$$



## » Punchline

Elements of  $\mathcal{M}$  correspond to filled diagrams like



shape  $\longleftrightarrow$  skew-commutative  $\theta$  factor

filling  $\longleftrightarrow$  commutative  $\times$  factor

Conjecture [SS24]/Theorem [ACK<sup>+</sup>24]

$$\mathcal{M} = \bigcup_{J \subseteq [n]} \{x^\alpha \theta_J : \alpha \leq (J\text{-staircase})\}$$

is a basis for  $SR_n$ .

What are SW arrangements and  $\mathcal{ST}$  algebras?

For the rest of the talk  $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ .



A hyperplane  $H$  is a codimension-1 affine linear subspace of  $\mathbb{K}^n$ .


A (hyperplane) arrangement  $\mathcal{A}$  is a union of hyperplanes.

Geometrically,  $H$  is a variety cut out by a degree-1 polynomial.

$\mathcal{A}$  is a variety cut out by a product of degree-1 polynomials.


## » Examples

An arrangement in  $\mathbb{R}^2$ :



“Definition”

Define by example the diagram  $\tilde{\Phi}_n$  for  $n = 5$ :




### Definition/example

An arrangement  $\mathcal{A}$  is called a *southwest arrangement* if its defining polynomial  $Q(\mathcal{A})$  is a product of terms of a southwest-closed subset of  $\tilde{\Phi}_n$ .


The  $h$ -function of a southwest arrangement is the number of hyperplanes on each southeast diagonal.

# » Example



is a southwest arrangement with  $h$ -function  $(1, 2)$ .

## » Example



$$x_1 x_2 (x_1 - x_2) (x_1 - x_3) (x_1 - x_4) (x_2 - x_3) (x_2 - x_4) (x_2 - x_5) (x_3 - x_4)$$

is southwest. It has  $h$ -function  $(1, 2, 2, 3, 1)$ .

There are myriad algebraic tools available to study arrangements.

### Definition

There is a free  $S$ -module  $\text{Der}(S)$  with basis  $\{\partial_i\}_{i=1}^n$ . The module of derivations of  $\mathcal{A}$  is

$$\text{Der}(\mathcal{A}) = \{d \in \text{Der}(S) : d(H) \in \text{span}_S H \forall H \in \mathcal{A}\}.$$

An arrangement is called free if  $\text{Der}(\mathcal{A})$  is a free  $S$ -module.

### Example

The arrangement  $x_1x_2(x_2 - x_1)$  is free with basis

$$\{x_1\partial_1 + x_2\partial_2, x_2(x_2 - x_1)\partial_2\}$$

If  $\mathcal{A}$  is a free arrangement, then  $\text{Der}(\mathcal{A})$  has a homogeneous basis.

The degrees of polynomials in the homogeneous basis are called the exponents of  $\mathcal{A}$ .

Exponents are independent of choice of basis and give combinatorial information about  $\mathcal{A}$ .



Theorem [ACK<sup>+</sup>24]

Let  $\mathcal{A}$  be a southwest arrangement. Then  $\mathcal{A}$  is free with exponents given by the  $h$ -function.

Definition [AMMN19]

Let  $\alpha : \text{Der}(\mathcal{A}) \rightarrow S$  be an  $S$ -module homomorphism.  
Define the Solomon-Terao algebra to be

$$ST(\mathcal{A}; \alpha) = S / \text{im } \alpha.$$

## » Example

Define  $\mathfrak{i}$  to be the map that sends  $\partial_i \mapsto 1$ .

Recall that the arrangement  $\mathcal{A}$  defined by  $x_1x_2(x_2 - x_1)$  is free with basis

$$\{x_1\partial_1 + x_2\partial_2, x_2(x_2 - x_1)\partial_2\}.$$

Then

$$ST(\mathcal{A}, \mathfrak{i}) = \mathbb{C}[x_1, x_2]/(x_1 + x_2, x_2^2 - x_1x_2) \cong \mathbb{C}[x_2]/(x_2^2).$$

Proof ideas

The colon ideal  $(I : f)$  is the kernel of  $\times f$  so that

$$0 \rightarrow S/(I : f) \xrightarrow{\times f} S/I$$

is exact.

Geometrically, for a variety  $X$ ,

$$(I(X) : f) = I(X - V(f))$$

## » Transfer principle

Rhoades and Wilson [RW23] showed that in order to show  $\mathcal{M}$  is a basis for  $SR_n$ , it suffices to show

$$\mathcal{M}(J) = \{x^\alpha : \alpha \leq (J\text{-staircase})\}$$

is a basis for

$$S/(I^+ : f_J)$$


where

$$f_J = \prod_{j \in J} x_j \prod_{i > j} (x_j - x_i).$$

### Upshot

Trade a skew-commutative problem for a family of commutative problems.

The arrangement defined by  $f_J$  is not a southwest arrangement.  
 E.g.  $J = \{2, 3\}$ :



### Theorem [ACK<sup>+</sup>24]

Let  $\mathcal{A}$  be an essential southwest arrangement in  $\mathbb{C}^n$  with  $h$ -function  $h(\mathcal{A})$ . Let  $i : \text{Der}(\mathcal{A}) \rightarrow S$  be defined by  $\partial_i \mapsto 1$ . Then the monomials

$$\{x^\alpha : \alpha < h(\mathcal{A})\}$$

descend to a basis for  $ST(\mathcal{A}; i)$ .

### Example

We saw  $\{1, x_2\}$  is a basis for  $ST(\mathcal{A}; i)$  of





### Definition

Let  $J \subseteq [n]$ . Let  $\mathcal{A}_J$  denote the southwest arrangement defined by

$$x_1 x_2 \cdots x_n \prod_{j \notin J} \prod_{i > j} (x_j - x_i)$$

### Example

Let  $J = \{2, 4\}$ , then  $\mathcal{A}_J$  is



### Definition

Let

$$\tilde{f}_J = \prod_{j \in J} \prod_{i > j} (x_j - x_i)$$

This is almost  $f_J$  but without the monomial factors.



Lemma [ACK<sup>+</sup>24]

$$S/(I_n^+ : \tilde{f}_J) \cong ST(\mathcal{A}_J; t)$$

Lemma

The  $J$ -staircase is bounded above by the  $h$ -function of  $\mathcal{A}_J$ . In particular

$$h_k = \begin{cases} \text{st}(J)_k & k \notin J \\ \text{st}(J)_k + 1 & k \in J \end{cases}$$



Theorem [ACK<sup>+</sup>24]

$\mathcal{M}(J)$  is a basis for  $S/(I^+ : f_J)$

Proof: A general fact about colon ideals tells us

$$(I^+ : \tilde{f}_J) : \underline{x}^J = (I^+ : \tilde{f}_J \underline{x}^J) = (I^+ : f_J).$$

Thus,

$$0 \rightarrow S/(I^+ : f_J) \xrightarrow{\times \underline{x}^J} S/(I^+ : \tilde{f}_J) = \mathcal{ST}(\mathcal{A}_J, \mathfrak{i})$$

is exact.





Corollary

$\mathcal{M}$  is a basis for  $SR_n$ , resolving conjecture of [SS24]

What next?

THANK YOU!

## » References

-  Robert Angarone, Patricia Commins, Trevor Karn, Satoshi Murai, and Brendon Rhoades, Superspace coinvariants and hyperplane arrangements, 2024.
-  Takuro Abe, Toshiaki Maeno, Satoshi Murai, and Yasuhide Numata, Solomon-Terao algebra of hyperplane arrangements, J. Math. Soc. Japan **71** (2019), no. 4, 1027–1047. MR 4023295
-  Brendon Rhoades and Andy Wilson, The hilbert series of the superspace coinvariant ring, 2023.
-  Bruce E. Sagan and Joshua P. Swanson,  $q$ -Stirling numbers in type  $B$ , European J. Combin. **118** (2024), Paper No. 103899, 35. MR 4674564

In case Sarah asks “What about type- $B$ ?”:

