

Equivariant Kazhdan–Lusztig theory of paving matroids

by Trevor K. Karn (U. Minnesota)

(joint with George Nasr, Nick Proudfoot, and Lorenzo Vecchi)


on Friday, February 17, 2023

A classical story

Our story

The nitty-gritty


Proof ideas



A partition $\lambda \vdash n$ is a weakly decreasing sequence of nonnegative integers $\lambda_1 \geq \lambda_2 \geq \dots$ summing to n .

Example

$\lambda = (5, 4, 4, 2, 1) \vdash 16$ has Ferrers diagram



A Young tableau T is a filling of a Ferrers diagram by positive integers. T is standard if it is filled by $\{1, 2, \dots, n\}$ and increasing in rows and columns. Define f^λ as the number of standard tableaux of shape λ .

Example

One of $f^{(5,4,4,2,1)} = 549120$ standard Young tableaux:

1	6	10	13	16
2	7	11	14	
3	8	12	15	
4	9			
5				

Fact

Fix n . Then

$$\sum_{\lambda \vdash n} (f^\lambda)^2 = n!$$

Proof 1:

The Robinson-Schensted bijection:pairs of standard tableaux of same shape \longleftrightarrow symmetric group \mathfrak{S}_n

Fact

The Specht modules S^λ are irreducible \mathfrak{S}_n representations indexed by $\lambda \vdash n$ and

$$\dim S^\lambda = f^\lambda.$$

Fact

Let d_1, d_2, \dots, d_r be the dimensions of the irreducible complex representations of a finite group. Then

$$\sum_i d_i^2 = |G|.$$


Fact

Fix n . Then

$$\sum_{\lambda \vdash n} (f^\lambda)^2 = n!$$

Proof 2:


$$\sum_{\lambda} (f^\lambda)^2 = \sum_i d_i^2 = |G| = |\mathfrak{S}_n| = n!$$



A skew partition λ/μ is a pair of partitions where the diagram of μ is contained in the diagram of λ

Example

If $\lambda = (4, 3, 2, 1)$ and $\mu = (2, 1)$ then λ/μ has diagram



A skew tableau T is a filling of a skew diagram by positive integers. T is standard if it is filled by $\{1, 2, \dots, |\lambda| - |\mu|\}$ and increasing in rows and columns. Define $f^{\lambda/\mu}$ as the number of standard skew tableaux of shape λ/μ .

Example

Two of $f^{(4,3,2,1)/(2,1)} = 272$ standard skew tableaux:

		1	2
	3	4	
5	6		
7			

		3	7
	2	6	
1	5		
4			

Theorem (Lee, Nasr, Radcliffe '21)



Let $P_M(t)$ be the matroid Kazhdan–Lusztig polynomial of M , a rank- k , sparse paving matroid with groundset $[n]$ and circuit hyperplanes \mathcal{CH} . The t^i coefficient in $P_M(t)$ is

$$f^{\lambda/\mu} - |\mathcal{CH}| f^{\lambda'/\mu'}$$

where

$$\lambda = [n - 2i, (k - 2i + 1)^i], \mu = [(k - 2i - 1)^i]$$

$$\lambda' = [(k - 2i + 1)^{i+1}], \mu' = [k - 2i, (k - 2i - 1)^{i-1}]$$



Theorem (Lee, Nasr, Radcliffe '21)

Let $P_M(t)$ be the matroid Kazhdan–Lusztig polynomial of M , a rank- k , sparse paving matroid with groundset $[n]$ and circuit hyperplanes \mathcal{CH} . The t^i coefficient in $P_M(t)$ is

$$f^{\lambda/\mu} - |\mathcal{CH}| f^{\lambda'/\mu'}$$

where

$$\lambda = [n - 2i, (k - 2i + 1)^i], \mu = [(k - 2i - 1)^i]$$

$$\lambda' = [(k - 2i + 1)^{i+1}], \mu' = [k - 2i, (k - 2i - 1)^{i-1}]$$

Proof 1 (LNR '21): Combinatorial argument with recursion.

Definition

The skew Specht module $S^{\lambda/\mu}$ is

$$S^{\lambda/\mu} = \bigoplus_{\nu} (S^{\nu})^{\oplus c_{\mu,\nu}^{\lambda}}$$

where $c_{\mu,\nu}^{\lambda}$ are Littlewood–Richardson coefficients.

Fact

$S^{\lambda/\mu}$ are (reducible) \mathfrak{S}_n representations and

$$\dim S^{\lambda/\mu} = f^{\lambda/\mu}.$$

Theorem (Lee, Nasr, Radcliffe '21)

Let $P_M(t)$ be the matroid Kazhdan–Lusztig polynomial of M , a rank- k , sparse paving matroid with groundset $[n]$ and circuit hyperplanes \mathcal{CH} . The t^i coefficient in $P_M(t)$ is

$$f^{\lambda/\mu} - |\mathcal{CH}| f^{\lambda'/\mu'}$$

where

$$\lambda = [n - 2i, (k - 2i + 1)^i], \mu = [(k - 2i - 1)^i]$$

$$\lambda' = [(k - 2i + 1)^{i+1}], \mu' = [k - 2i, (k - 2i - 1)^{i-1}]$$

Proof 1 (LNR '21): Combinatorial argument with recursion.

Proof 2 (KNPV '22): $\dim(\text{skew Specht module from } M)$.

» Example: $U_{3,12}$




$$\mathcal{CH} = \emptyset$$


$$f^{(10,2)} = 54$$

$$P_{U_{3,12}}(t) = 1 + 54t$$

» Example: Vamos matroid



$$\lambda = [6, 3], \mu = [1] \longrightarrow$$


$$\lambda' = [3, 3], \mu' = [2] \longrightarrow$$


$$|\mathcal{CH}| = 5$$

$$f^{\lambda/\mu} - 5f^{\lambda'/\mu'} = 48 - 15 = 33$$

$$P_V(t) = 1 + 33t$$

» Example: Projective plane over \mathbb{F}_3



$$C\mathcal{H} = \emptyset$$

$$|S\mathcal{H}| = 13$$



$$f^{\lambda/\mu} - 13f^{\lambda'/\mu'} = 65 - 13 * 5 = \neq 0$$

$P_M(t) = 1$

Theorem

Let $P_M(t)$ be the matroid Kazhdan–Lusztig polynomial of M , a rank- k , (arbitrary!) paving matroid with groundset $[n]$ and nontrivial stressed hyperplanes \mathcal{SH} . The t^i coefficient in $P_M(t)$ is



$$f^{\lambda/\mu} - \sum_{H \in \mathcal{SH}} f^{\lambda'(H)/\mu'}$$

where

$$\lambda = [n - 2i, (k - 2i + 1)^i], \mu = [(k - 2i - 1)^i]$$

$$\lambda'(h) = [h - 2i + 1, (k - 2i + 1)^i], \mu' = [h - 2i, (k - 2i - 1)^{i-1}]$$

Proof: Our proof of LNR's theorem implies this more general result



Certain $P_M(t)$ in terms of standard skew Young tableaux



Combinatorial proof [LNR21]

Algebraic proof [KNPV22]

» What do I owe you?

Matroids

Circuits and stressed hyperplanes
(Sparse) paving

Kazhdan–Lusztig polynomials

How $S^{\lambda/\mu}$ arises


“Definition” 1

A matroid $M = (E, \mathcal{B})$ is a finite set E (called the **groundset**) together with $\mathcal{B} \subseteq 2^E$ satisfying some axioms combinatorially modeling choices of bases for a vector space.

Alternatively...

“Definition” 2

A matroid $M = (E, \mathcal{C})$ is a ground set E together with $\mathcal{C} \subseteq 2^E$ satisfying some axioms modeling minimal linear dependence of vectors.



Bases \longleftrightarrow maximal independent sets

Circuits \longleftrightarrow minimal dependent sets

Example

The **uniform matroid** $U_{k,n}$ models n -many k -dimensional vectors in general position


Bases \longleftrightarrow any set of k -many vectors

Circuits \longleftrightarrow any set of $k + 1$ -many vectors

Example of the example

$U_{3,12}$ corresponds to 12 generic vectors in \mathbb{R}^3 . One choice of basis is $\{e_1, e_2, e_3\}$. On the other hand $\{e_1, e_2, e_3, v\}$ is dependent for any $v \in \mathbb{R}^3$.

The combinatorial model for: vectors \rightarrow groundset elements
 subspaces \rightarrow flats




Flats form a ranked lattice L . Define $r(M) = r(L) = k$.
 Rank- $(k - 1)$ flats are **hyperplanes**. A **circuit hyperplane** is also
 a circuit.

M is a paving matroid if all circuits are at least size $k = r(M)$

A paving matroid is sparse if the set \mathcal{CH} of circuit hyperplanes satisfies $\binom{E}{k} = \mathcal{CH} \sqcup \mathcal{B}$

A circuit hyperplane is the prototypical example of...

a stressed hyperplane H of a rank- k matroid has every k -subset a circuit.



Conjecture (Mayhew, Newman, Welsh, Whittle '11)

Asymptotically almost all matroids are sparse paving
(\Rightarrow paving)

Theorem (Pendavingh, van der Pol '15)

Asymptotically logarithmically almost all matroids are
sparse paving

» What do I owe you?

Matroids ✓

 Circuits and stressed hyperplanes ✓

 (Sparse) paving ✓

Kazhdan–Lusztig polynomials

How $S^{\lambda/\mu}$ arises

In order to define P_M , first define

$$\chi_M(t) = \sum_{F \in L(M)} \mu(\bar{\emptyset}, F) t^{k-r(F)}$$

where μ is the Möbius function.

Definition/Theorem (Elias, Proudfoot, Wakefield '16)


Fix M . There exists a unique polynomial $P_M(t)$ satisfying:

$$P_M(t) = 1 \text{ if } r(M) = 0,$$

$$\deg P_M(t) < r(M)/2 \text{ when } r(M) > 0,$$

$$t^{r(M)} \overline{P_M}(t) = \sum_{F \in L(M)} P_{M_F}(t) \chi_{M^F}(t).$$

M_F and M^F



» What do I owe you?

Matroids ✓

 Circuits and stressed hyperplanes ✓

 (Sparse) paving ✓

Kazhdan–Lusztig polynomials ✓

How $S^{\lambda/\mu}$ arises

Let W be a group. An equivariant matroid $W \curvearrowright M$ is a matroid with a W -action “preserving the matroid.”

e.g. $gB \in \mathcal{B}$ for all $g \in W$ and $B \in \mathcal{B}$


gF is another flat of the same rank

The Orlik–Solomon algebra $\mathcal{OS}(M)$ is a certain quotient of the exterior algebra $\bigwedge^\bullet K^n$

Theorem (Orlik, Solomon '80)

$\chi_M(t)$ determines the Poincaré polynomial of $\mathcal{OS}(M)$

$W \curvearrowright M$ induces a W -action on $\mathcal{OS}(M)$. Use this to define a graded virtual representation called the equivariant characteristic polynomial. The coefficient of t^{k-i} is $\pm \mathcal{OS}(M)_i$.



Definition/Theorem (Gedeon, Proudfoot, Young '17)

Let $W \curvearrowright M$ be an equivariant matroid. Then there exists a unique $P_M^W(t)$ with

If $r(M) = 0$, then $P_M^W(t)$ is $\mathbb{1}_W t^0$

If $r(M) > 0$, then $\deg P_M^W(t) < r(M)/2$

$$t^{r(M)} \overline{P}_M^W(t) = \sum_{[F] \in L(M)/W} \text{Ind}_{W_F}^W \left(P_{M_F}^{W_F}(t) \otimes \chi_{M_F}^{W_F} \right)$$

$\varphi : W' \rightarrow W$ a homom. then $P_M^{W'}(t) = \varphi^* P_M^W(t)$

where W_F denotes the stabilizer of F .

Compare:

$$t^{r(M)} \overline{P}_M(t) = \sum_{F \in L(M)} P_{M_F}(t) \chi_{M^F}(t)$$

and


$$t^{r(M)} \overline{P}_M^W(t) = \sum_{[F] \in L(M)/W} \text{Ind}_{W_F}^W \left(P_{M_F}^{W_F}(t) \otimes \chi_{M^F}^{W_F} \right).$$




Theorem (Ferroni, Nasr, Vecchi '21)

Let $M = (E, \mathcal{B})$ be a matroid with stressed hyperplane H . The operation of relaxation at H forms a new matroid $\tilde{M} = (E, \tilde{\mathcal{B}})$ with bases

$$\tilde{\mathcal{B}} = \mathcal{B} \sqcup \{S \subseteq H : |S| = k\}.$$





Theorem (Ferroni, Nasr, Vecchi '21)

There exists a polynomial $p_{k,h}$ such that

$$P_M(t) = P_{\tilde{M}}(t) - p_{k,h}$$

Fact


M is paving \Leftrightarrow a sequence of relaxations makes it $U_{k,n}$

Theorem (Ferroni, Nasr, Vecchi '21)

If M is a paving matroid with $|E| = n$ and has exactly λ_h -many stressed hyperplanes of size h , then

$$P_M(t) = P_{U_{k,n}}(t) - \sum_{h \geq k} \lambda_h \cdot p_{k,h}.$$

» Idea of the proof



Let $W \curvearrowright M$ be an equivariant matroid with stressed hyperplane H .

Let $W \curvearrowright \tilde{M}$ denote the equivariant matroid found by simultaneously relaxing all hyperplanes in $[H]$.

Theorem (K.-Nasr-Proudfoot-Vecchi '22)

There exists an equivariant polynomial $p_{k,h}^{\mathfrak{S}_h}$ such that

$$P_M^W(t) = P_{\tilde{M}}^W(t) - \text{Ind}_{W_H}^W \text{Res}_{W_H}^{\mathfrak{S}_h} p_{k,h}^{\mathfrak{S}_h}$$

Theorem (K.-Nasr-Proudfoot-Vecchi '22)



The coefficients of t^i are

$$\{t^i\} p_{k,h}^{\mathfrak{S}_h} = S^{\lambda'/\mu'}$$

where $\lambda', \mu' \vdash h$ are:

$$\lambda' = h - 2i + 1, (k - 2i + 1)^i \text{ and}$$

$$\mu' = k - 2i, (k - 2i - 1)^{i-1}$$




» Idea of proof

Relax $U_{k-1,h}^{\mathfrak{S}_h} \oplus U_{1,1}$ to $U_{k,h+1}^{\mathfrak{S}_{h+1}}$.


$$P_{M_1 \oplus M_2}(t) = P_{M_1}(t)P_{M_2}(t)$$

$P_{k,h}^{\mathfrak{S}_h}$ depends only on k, h , so one example is enough.



Theorem (Gao, Xie, Yang '21)

Every coefficient of t^i in $P_{U_{k,n}}^{\mathfrak{S}_n}(t)$ is given by the skew Specht module of shape



Combine:

M is paving \Leftrightarrow a sequence of relaxations makes it $U_{k,n}$

Theorems (K.-Nasr-Proudfont-Vecchi '22)

$$P_M^W(t) = P_{\tilde{M}}^W(t) - \text{Ind}_{W_H}^W \text{Res}_{W_H}^{\mathfrak{S}_h} p_{k,h}^{\mathfrak{S}_h}$$

and coefficients of $p_{k,h}^{\mathfrak{S}_h}$ are $S^{\lambda(h)/\mu}$

Theorem (Gao, Xie, Yang '21)

Coefficients of $P_{U_{k,n}}^{\mathfrak{S}_n}(t)$ are $S^{\lambda/\mu}$

$$\dim(S^{\lambda/\mu}) = f^{\lambda/\mu}$$



to obtain...

Theorem

Let $P_M(t)$ be the matroid Kazhdan–Lusztig polynomial of M , a rank- k , arbitrary paving matroid with groundset $[n]$ and nontrivial stressed hyperplanes \mathcal{SH} . The t^i coefficient in $P_M(t)$ is

$$f^{\lambda/\mu} - \sum_{H \in \mathcal{SH}} f^{\lambda'(|H|)/\mu'}$$

where λ/μ , λ'/μ' are as before.



THANK YOU!

» References



Ben Elias, Nicholas Proudfoot, and Max Wakefield, The Kazhdan–Lusztig polynomial of a matroid, *Advances in Mathematics* **299** (2016), 36–70.



Luisa Ferroni, George D. Nasr, and Lorenzo Vecchi, Stressed hyperplanes and Kazhdan–Lusztig gamma-positivity for matroids, 2021.



Katie R. Gedeon, Nicholas Proudfoot, and Benjamin Young, The equivariant Kazhdan–Lusztig polynomial of a matroid, *J. Comb. Theory, Ser. A* **150** (2017), 267–294.



Alice L. L. Gao and Matthew H. Y. Xie, The inverse Kazhdan–Lusztig polynomial of a matroid, *J. Combin. Theory Ser. B* **151** (2021), 375–392. MR 4294228

» References (cont.)



Alice L. L. Gao, Matthew H.Y. Xie, and Arthur L. B. Yang, The equivariant inverse Kazhdan–Lusztig polynomials of uniform matroids, 2021.



Trevor Karn, George Nasr, Nicholas Proudfoot, and Lorenzo Vecchi, Equivariant Kazhdan–Lusztig theory of paving matroids, 2022.



Kyungyong Lee, George D. Nasr, and Jamie Radcliffe, A combinatorial formula for Kazhdan–Lusztig polynomials of sparse paving matroids, *Electron. J. Comb.* **28** (2021).



Dillon Mayhew, Mike Newman, Dominic Welsh, and Geoff Whittle, On the asymptotic proportion of connected matroids, *European J. Combin.* **32** (2011), no. 6, 882–890. MR 2821559

» References (cont.)



Peter Orlik and Louis Solomon, Combinatorics and topology of complements of hyperplanes, *Inventiones mathematicae* **56** (1980), no. 2, 167–189 (eng).



Rudi Pendavingh and Jorn van der Pol, On the number of matroids compared to the number of sparse paving matroids, *Electron. J. Combin.* **22** (2015), no. 2, Paper 2.51, 17. MR 3367294



Nicholas Proudfoot, Yuan Xu, and Benjamin Young, The Z-polynomial of a matroid, *Electron. J. Comb.* **25** (2018), P1.26.